



## Averting the legacy of kidney disease—focus on childhood

Julie R. Ingelfinger<sup>1</sup>, Kamyar Kalantar-Zadeh<sup>2</sup> and Franz Schaefer<sup>3</sup>; for the World Kidney Day Steering Committee<sup>4</sup>  
<sup>1</sup>Professor of Pediatrics, Harvard Medical School, Senior Consultant in Pediatric Nephrology and Pediatric Nephrology and Pediatric, Mass General Hospital for Children at Massachusetts General Hospital, Boston, Massachusetts, USA; <sup>2</sup>Chief, Division of Nephrology & Hypertension, Professor of Medicine, Pediatrics and Public Health, University of California Irvine, School of Medicine; Staff Physician and Principal Investigator, Veterans Affairs (VA) Long Beach Healthcare System, Long Beach, CA, USA; Adjunct Professor of Epidemiology, Department of Epidemiology, UCLA Fielding School of Public Health, Los Angeles, CA, USA; and <sup>3</sup>Professor of Pediatrics, University of Heidelberg, Head, Division of Pediatric Nephrology, Center for Pediatrics and Adolescent Medicine Heidelberg, Germany  
**Correspondence:** World Kidney Day, International Society of Nephrology, in collaboration with International Federation of Kidney Foundation, Rues de Fabriques 1B, 1000, Brussels, Belgium. E-mail: [myriam@worldkidneyday.org](mailto:myriam@worldkidneyday.org)

<sup>4</sup>See Appendix for list of committee members.

This article is being published in *Kidney International* and reprinted concurrently in several journals. The articles are identical except for minor stylistic and spelling changes in keeping with each journal's style. Any of these versions may be used in citing this article.

World Kidney Day 2016 focuses on kidney disease in childhood and the antecedents of adult kidney disease that can begin in earliest childhood. Chronic kidney disease in childhood differs from that in adults, as the largest diagnostic group among children includes congenital anomalies and inherited disorders, with glomerulopathies and kidney disease in the setting of diabetes being relatively uncommon. In addition, many children with acute kidney injury will ultimately develop sequelae that may lead to hypertension and chronic kidney disease in later childhood or in adult life. Children born early or who are small-for-date newborns have a relatively increased risk for the development of chronic kidney disease later in life. Persons with a high-risk birth and early childhood history should be watched closely in order to help detect early signs of kidney disease in time to provide effective prevention or treatment. Successful therapy is feasible for advanced chronic kidney disease in childhood; there is evidence that children fare better than adults if they receive kidney replacement therapy including dialysis and transplant, whereas only a minority of children may require this ultimate intervention. Because there are disparities in access to care, effort is needed so that those children with kidney disease, wherever they live, may be treated effectively, irrespective of their geographic or economic circumstances. Our hope is that World Kidney Day will inform the general public, policy makers, and caregivers about the needs and possibilities surrounding kidney disease in childhood.

*Kidney International* (2016) **89**, 512–518; <http://dx.doi.org/10.1016/j.kint.2015.10.014>  
 © World Kidney Day 2016 Steering Committee

“For in every adult there dwells the child that was, and in every child there lies the adult that will be.”


—John Connolly, *The Book of Lost Things*

The 11<sup>th</sup> World Kidney Day will be celebrated on March 10, 2016, around the globe. This annual event, sponsored jointly by the International Society of Nephrology (ISN) and the International Federation of Kidney Foundations, has become a highly successful effort to inform the general public and policy makers about the importance and ramifications of kidney disease. In 2016, World Kidney Day will be dedicated to kidney disease in childhood and the antecedents of adult kidney disease, which can begin in earliest childhood.

Children who endure acute kidney injury (AKI) from a wide variety of conditions may have long-term sequelae that can lead to chronic kidney disease (CKD) many years later.<sup>1–4</sup> Furthermore, CKD in childhood, much of it congenital, and complications from the many nonrenal diseases that can affect the kidneys secondarily, not only lead to substantial morbidity and mortality during childhood, but also result in medical issues beyond childhood (Figure 1). Indeed, childhood deaths from a

long list of communicable diseases are inextricably linked to kidney involvement. For example, children who succumb to cholera and other diarrheal infections often die, not from the infection, but because of AKI induced by volume depletion and shock. In addition, a substantial body of data indicates that hypertension, proteinuria, and CKD in adulthood have childhood antecedents—from as early as *in utero* and perinatal life (see Table 1 for definitions of childhood). World Kidney Day 2016 aims to heighten general awareness that much adult renal disease is actually initiated in childhood. Understanding high-risk diagnoses and events that occur in childhood have the potential to identify and intervene preemptively in those people at higher risk for CKD during their lifetimes.

Worldwide epidemiologic data on the spectrum of both CKD and AKI in children are currently limited, though increasing in scope. The prevalence of CKD in childhood is rare—and has been variously reported at 15 to 74.7 per million children.<sup>3</sup> Such variation is likely because data on CKD are influenced by regional and cultural factors, as well as by the methodology used to generate them. The World Health Organization has recently added



**Figure 1 | The types and risks of kidney disease change across the life cycle.** The contribution of nephron number increases over the life cycle, in concert with events that provide direct insults and challenges to kidney health.

kidney and urologic disease to mortality information tracked worldwide and should be a valuable source of such data over time, but the World Health Organization does not post the information by age group.<sup>5</sup> Databases such as the North American Pediatric Renal Trials and Collaborative Studies (NAPRTCS)<sup>6</sup> the U.S. Renal Data System (USRDS)<sup>7</sup> and the European Dialysis and Transplant Association (EDTA) registry<sup>8</sup> include data on pediatric end-stage renal disease (ESRD), and some on CKD. Projects such as the ItalKid<sup>9</sup> and Chronic Kidney Disease in Children (CKiD)<sup>10</sup> studies, the Global Burden of Disease Study 2013, as well as registries that now exist in many countries provide important information, and more is required.<sup>11</sup>

AKI may lead to CKD, according to selected adult population studies.<sup>12</sup> The incidence of AKI among children admitted to an intensive care unit varies widely—from 8% to 89%.<sup>1</sup> The outcome depends on the available resources. The results from projects such as the

Assessment of Worldwide Acute Kidney Injury, Renal Angina and Epidemiology (AWARE) study, a 5-nation study of AKI in children are awaited.<sup>13</sup> Single-center studies, as well as meta-analyses indicate that both AKI and CKD in children account for a minority of CKD worldwide.<sup>2,3</sup> However, it is increasingly evident that kidney disease in adulthood often springs from a childhood legacy.

**SPECTRUM OF PEDIATRIC KIDNEY DISEASES**

The conditions that account for CKD in childhood, with a predominance of congenital and hereditary disorders, differ substantially from those in adults. To date, mutations in >150 genes have been found to alter kidney development or specific glomerular or tubular functions.<sup>14</sup> Most of these genetic disorders present during childhood, and many lead to progressive CKD. Congenital anomalies of the kidney and urinary tract account for the largest category of CKD in children (see Table 2) and include renal hypoplasia/dysplasia and obstructive uropathy. Important subgroups among the renal dysplasias are the cystic kidney diseases, which originate from genetic defects of the tubuloepithelial cells’ primary cilia. Many pediatric glomerulopathies are caused by genetic or acquired defects of the podocytes, the unique cell type lining the glomerular capillaries. Less common but important causes of childhood CKD are inherited metabolic disorders such as hyperoxaluria and cystinosis, and atypical hemolytic uremic syndrome, a thrombotic microangiopathy related to genetic abnormalities of complement, coagulation, or metabolic pathways.

In various classifications, it is not clear how to categorize children who have suffered AKI

**Table 1 | Definitions of stages of early life**

Perinatal period	22 Completed weeks of gestation to day 7 of postnatal life
Neonatal period	Birth to day 28 of postnatal life
Infancy	Birth to 1 year of age
Childhood	1 Year of age to 10 years of age
Adolescence	10 Years of age to 19 years of age

The data in this table are as defined by the World Health Organization. The perinatal period is defined as 22 completed weeks of gestation to day 7 of life; the neonatal period as up to 28 days of life; infancy as up to 1 year of age; childhood as years 1 to 10; and adolescence from 10 years to age 19.

There is variation worldwide in how these stages of early life are defined. Some would define “young people” as those age 24 or younger. In the United States, childhood is as a whole defined as going to age 21.

**Table 2 | Etiology of chronic kidney disease in children**

CKD etiology	Percentage range	ESRD etiology	Percentage range
CAKUT	48–59	CAKUT	34–43
GN	5–14	GN	15–29
HN	10–19	HN	12–22
HUS	2–6	HUS	2–6
Cystic	5–9	Cystic	6–12
Ischemic	2–4	Ischemic	2

CAKUT, congenital anomalies of the kidney and urinary tract; CKD, chronic kidney disease; ESRD, end-stage renal disease; GN, glomerulonephritis; HN, hypertension; HUS, hemolytic uremic syndrome. Rare causes include congenital nephrotic syndrome, metabolic diseases, and cystinosis. Miscellaneous causes depend on how such entities are classified.

Adapted from Harambat et al.<sup>2</sup> CKD data are from North American Pediatric Renal Trials and Collaborative Studies NAPRTCS, the Italian Registry, and the Belgian Registry. ESRD data are from Australia and New Zealand Dialysis and Transplant Registry (ANZDATA), European Society of Pediatric Nephrology/European Renal Association–European Dialysis and Transplant Association (ESPN/ERA-EDTA) Registry, U.K. Renal Registry, and the Japanese Registry.

and apparently recovered, or how and whether to include those children who have had perinatal challenges, likely resulting in a relatively low nephron number.

Among children with childhood-onset ESRD, glomerulopathies are slightly more and congenital anomalies less common (Table 2) due to the typically more rapid nephron loss in glomerular disease. However, recent evidence suggests that many patients with milder forms of congenital anomalies of the kidney and urinary tract may progress to ESRD during adulthood, peaking in the fourth decade of life.<sup>15</sup>

There are national and regional differences in the types and course of both AKI and CKD during childhood and beyond. Death from kidney disease is higher in developing nations, and national and regional disparities in care and outcome must be addressed. Furthermore, access to care is variable, depending on the region, the country and its infrastructure. By focusing on kidney disease in childhood, cost-effective solutions may be reached, as treating disease early and pre-emptively may prevent later, more advanced CKD. Expectations depend on the availability of care and management. Treating children, even from infancy, who have AKI and CKD that requires renal replacement therapy (RRT) can be effective in mitigating the burden of kidney disease in adulthood. Doing so requires resources that focus on the most expeditious and cost-effective ways to deliver acute RRT in childhood.

**CONGENITAL KIDNEY DISEASE AND DEVELOPMENTAL ORIGINS OF HEALTH AND DISEASE, RENAL ENDOWMENT, AND IMPLICATIONS**

In regions where antenatal fetal ultrasonograms are routine, many children with urologic

abnormalities are identified antenatally, which permits early intervention. However, in much of the world, children with structural abnormalities are not identified until much later, when symptoms develop. Although generalized screening for proteinuria, hematuria, and urinary tract infections are carried out in some countries and regions, there is a lack of consensus as to its effectiveness. However, there is general agreement that children with antenatal ultrasonogram studies that indicate possible genitourinary anomalies; children with a family history of kidney disease; and children with signs such as failure to thrive or a history of urinary tract infection, voiding dysfunction, or an abnormal appearing urine should be examined. Initial screening would include a focused physical examination and a urine dipstick, formal urinalysis, and a basic chemistry panel, followed by a more focused evaluation if indicated.

Depending on the diagnosis, definitive therapy may be indicated. However, the evidence that therapy will slow progression of CKD in childhood remains limited. Angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, antioxidants, and, possibly, dietary changes may be indicated, depending on the diagnosis. However, dietary changes need to permit adequate growth and development. The Evaluation Study of Congestive Heart Failure and Pulmonary Artery Catheterization Effectiveness (ESCAPE) trial provided evidence that strict blood pressure control retards progression of CKD in children irrespective of the type of underlying kidney disease.<sup>16</sup>

Some very young children may require RRT in early infancy. Recent data pooled from registries worldwide indicate good survival, even when dialysis is required from neonatal age.<sup>2,17</sup> Kidney transplantation, the preferred RRT in

children, is generally suitable after 12 months of age, with excellent patient and allograft survival, growth, and development.

Evidence is accumulating that childhood-onset CKD leads to accelerated cardiovascular morbidity and shortened life expectancy. Ongoing large prospective studies such as the Cardiovascular Comorbidity in Children with CKD (4C) study are expected to inform about the causes and consequences of early cardiovascular disease in children with CKD.<sup>18</sup>

In addition to those children with congenital kidney disease, it is now known that perinatal events may affect future health in the absence of evident kidney disease in early life.<sup>19</sup> Premature infants appear to be particularly at risk for kidney disease long after they are born, based both on observational cohort studies, as well as on case reports. Increasingly premature infants survive, including many born well before nephrogenesis is complete.<sup>20</sup> The limited data available indicate that in the process of neonatal intensive care unit care, such babies receive many nephrotoxins, and that those dying prior to discharge from the nursery have fewer and larger glomeruli.<sup>21</sup> Additionally, those surviving have evidence of renal impairment that may be subtle.<sup>22</sup> Even more concerning, abundant epidemiologic data indicate that persons born at term but with relatively low birth weights may be at high risk for hypertension, albuminuria, and CKD in later life.<sup>23</sup> When direct measurements are pursued, such persons, as adults, may have fewer nephrons, thus a low cardiorenal endowment.

In focusing on children for World Kidney Day, we would note that it is key to follow kidney function and blood pressure throughout life in those persons born early or small-for-date. By doing so, and avoiding nephrotoxic medications throughout life, it may be possible to avert CKD in many people.

#### **RESOURCES AND THERAPEUTICS FOR CHILDREN—DIFFERENCES FROM THERAPEUTICS IN ADULTS**

Disparities exist in the availability of resources to treat AKI in children and young people; consequently, too many children and young adults in developing nations succumb if AKI occurs. To address the problem, the ISN has initiated the Saving Young Lives Project, which aims both to prevent AKI with prompt treatment of infection and/or delivery of appropriate fluid and electrolyte therapy, and to

treat AKI when it occurs. This ongoing project in Sub-Saharan Africa and South East Asia, in which 4 kidney foundations participate equally (International Pediatric Nephrology Association, ISN, International Society for Peritoneal Dialysis, Sustainable Kidney Care Foundation), focuses on establishing and maintaining centers for the care of AKI, including the provision of acute peritoneal dialysis. It links with the ISN's 0 by 25 project, which calls on members to ensure by 2025 that nobody dies from preventable and acute kidney injury.

In view of the preponderance of congenital and hereditary disorders, therapeutic resources for children with CKD have historically been limited to a few immunological conditions. Very recently, progress in drug development in concert with advances in genetic knowledge and diagnostic capabilities has begun to overcome the longstanding "therapeutic nihilism" in pediatric kidney disease. Atypical hemolytic uremic syndrome, long considered ominous, with a high likelihood of progression to ESRD and posttransplantation recurrence, has turned into a treatable condition—with the advent of a monoclonal antibody that specifically blocks C5 activation.<sup>24</sup> Another example is the use of vasopressin receptor antagonists to retard cyst growth and preserve kidney function in polycystic kidney disease.<sup>25</sup> First proven efficacious in adults with autosomal dominant polycystic kidney disease, therapy with vaptans holds promise also for the recessive form of the disease, which presents and often progresses to ESRD during childhood.

However, patient benefit from pharmacological research breakthroughs is jeopardized on a global scale by the enormous cost of some of the new therapeutic agents. The quest for affordable innovative therapies for rare diseases will be a key issue in pediatric nephrology in the years to come.

The identification of children likely to benefit from novel therapeutic approaches will be greatly facilitated by the development of clinical registries that inform about the natural disease course, including genotype-phenotype correlations. Apart from disease-specific databases, there is also a need for treatment-specific registries. These are particularly relevant in areas where clinical trials are difficult to perform due to small patient numbers and lacking industry interest, as well as for therapies

in need of global development or improvement. For instance, there is currently a large international gradient in the penetration and performance of pediatric dialysis and transplantation. Whereas pediatric patient and technique survival rates are excellent and even superior to those of adults in many industrialized countries, it is estimated that almost one-half of the world's childhood population is not offered chronic RRT at all. Providing access to RRT for all children will be a tremendous future challenge. To obtain reliable information on the demographics and outcomes of pediatric RRT, the International Pediatric Nephrology Association is about to launch a global population-based registry. If successful, the International Pediatric Nephrology Association RRT registry might become a role model for global data collection.

#### **TRANSITION FROM PEDIATRIC TO ADULT CARE**

Transition of care for adolescents with kidney disease into an adult setting is critical both for patients and their caregivers. Nonadherence is a too-frequent hallmark of transition from pediatric to adult care for young patients with chronic disease states.<sup>26–28</sup> Hence, considered steps combined with systematically defined procedures supported by validated pathways and credible guidelines must be in place to ensure successful outcomes.

In the process of change from pediatric to adult care, “transition,” which should occur gradually, must be distinguished from “transfer,” which is often an abrupt and mechanistic change in provider setting. Introducing the concept of transition should be pre-emptive, starting months to years prior to the targeted time, as children move into adolescence and adulthood. The ultimate goal is to foster a strong relationship and individualized plan in the new setting that allows the patient to feel comfortable enough to report nonadherence and other lapses in care.

A transition plan must recognize that the emotional maturity of children with kidney disease may differ widely. Assessment of the caregiver and the family structure as well as cultural, social, and financial factors at the time of transition are key, including a realistic assessment of caregiver burden.<sup>4</sup> The appropriate timing and format of transition may vary widely among different patients and in different settings; therefore, a flexible process without a set date and even without a delineated format may be preferred.

Importantly, transition may need to be slowed, paused, or even reversed temporarily during crises such as disease flares or progression, or if family or societal instability occurs. A recent joint consensus statement by the ISN and International Pediatric Nephrology Association proposed steps consistent with the points just outlined, aiming to enhance the transition of care in kidney disease in clinical practice.<sup>29,30</sup>

#### **CALL FOR GENERATING FURTHER INFORMATION AND ACTION**

Given vulnerabilities of children with kidney disease including impact on growth and development and future life as an adult, and given the much greater proportion of children in developing nations facing resource constraints educating everyone involved is imperative in order to realign communications and actions.<sup>31,32</sup> These efforts should foster regional and international collaborations and exchange of ideas among local kidney foundations, professional societies, other not-for-profit organizations, and states and governments, so as to help empower all stakeholders to improve the health, well-being, and quality of life of children with kidney diseases and to ensure their longevity into adulthood.

Until recently, however, the World Health Organization consensus statement on non-communicable diseases (NCDs) included cardiovascular disease, cancer, diabetes, and chronic respiratory disease, but not kidney disease.<sup>33,34</sup> Fortunately, due, in part, to a global campaign led by the ISN, the Political Declaration on NCDs from the U.N. Summit in 2011 mentioned kidney disease under Item 19.<sup>35</sup>

Increasing education and awareness about renal diseases in general and kidney disease in childhood in particular is consistent with the objectives of the World Health Organization to reduce mortality from NCD with a 10-year target population-level initiatives focusing on changes in lifestyle (including tobacco use reduction, salt intake control, dietary energy control, and alcohol intake reduction) and effective interventions (including blood pressure, cholesterol, and glycemic control). Heightened efforts are needed to realign and expand these multidisciplinary collaborations with more effective focus on early detection and management of kidney disease in children. Whereas the issues related to kidney disease may be overshadowed by other NCDs with

apparently larger public health implications such as diabetes, cancer, and cardiovascular diseases, our efforts should also increase education and awareness on such overlapping conditions as cardiorenal connections, the global nature of the CKD and ESRD as major NCDs, and the role of kidney disease as the multiplier disease and confounder for other NCDs. White papers including consensus articles and blueprint reviews by world class experts can serve to enhance these goals.<sup>36</sup>

## REFERENCES

- Goldstein SL. Acute kidney injury in children and its potential consequences in adulthood. *Blood Purif.* 2012;33:131–137.
- Harambat J, van Stralen KJ, Kim JJ, Tizard EJ. Epidemiology of chronic kidney disease in children. *Pediatr Nephrol.* 2012;27:363–373.
- Warady BA, Chadha V. Chronic kidney disease in children: the global perspective. *Pediatr Nephrol.* 2007;22:1999–2009.
- Furth SL, Cole SR, Moxey-Mims M, et al. Design and methods of the Chronic Kidney Disease in Children (CKiD) prospective cohort study. *Clin J Am Soc Nephrol.* 2006;1:1006–1015.
- World Health Organization. Health statistics and information systems: Estimates for 2000–2012. Available at: [http://www.who.int/healthinfo/global\\_burden\\_disease/estimates/en/index1.html](http://www.who.int/healthinfo/global_burden_disease/estimates/en/index1.html). Accessed 26 October 2015.
- North American Pediatric Renal Trials and Collaborative Studies. NAPRTCS Annual Reports. Available at: <https://web.emmes.com/study/ped/annlrept.html>. Accessed 25 October 2015.
- Saran R, Li Y, Robinson B, et al. US Renal Data System 2014 Annual Data Report: epidemiology of kidney disease in the United States. *Am J Kidney Dis.* 2015;66(suppl 1):Svii, S1–305.
- ESPN/ERA-EDTA Registry, European Registry for Children on Renal Replacement Therapy. Welcome to the website of the ESPN/ERA-EDTA Registry. Available at: <http://www.espn-reg.org/index.jsp>. Accessed 25 October 2015.
- Ardissino G, Dacco V, Testa S, et al., for the Italkid Project. Epidemiology of chronic renal failure in children: data from the Italkid project. *Pediatrics.* 2003;111:e382–e387.
- Wong CJ, Moxey-Mims M, Jerry-Fluker J, et al. CKiD (CKD in Children) prospective cohort study: a review of current findings. *Am J Kidney Dis.* 2012;60:1002–1011.
- Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015;386:743–800.
- Coca SG, Singanamala S, Parikh CR. Chronic kidney disease after acute kidney injury: a systematic review and meta-analysis. *Kidney Int.* 2012;81:442–448.
- Basu RK, Kaddourah A, Terrell T, et al., for the ppAKI Research Group. Assessment of Worldwide Acute Kidney Injury, Renal Angina and Epidemiology in critically ill children (AWARE): study protocol for a prospective observational study. *BMC Nephrol.* 2015;16:24.
- Eckardt KU, Coresh J, Devuyst O, et al. Evolving importance of kidney disease: from subspecialty to global health burden. *Lancet.* 2013;382:158–169.
- Wuhl E, van Stralen KJ, Verrina E, et al. Timing and outcome of renal replacement therapy in patients with congenital malformations of the kidney and urinary tract. *Clin J Am Soc Nephrol.* 2013;8:67–74.
- Wuhl E, Trivelli A, Picca S, et al., for the ESCAPE Trial Group. Strict blood-pressure control and progression of renal failure in children. *N Engl J Med.* 2009;361:1639–1650.
- van Stralen KJ, Borzych-Dużalka D, Hataya H, et al. Survival and clinical outcomes of children starting renal replacement therapy in the neonatal period. *Kidney Int.* 2014;86:168–174.
- Querfeld U, Anarat A, Bayazit AK, et al., for the 4C Study Group. The Cardiovascular Comorbidity in Children with Chronic Kidney Disease (4C) study: objectives, design, and methodology. *Clin J Am Soc Nephrol.* 2010;5:1642–1648.
- Hoy WE, Ingelfinger JR, Hallan S, et al. The early development of the kidney and implications for future health. *J Dev Orig Health Dis.* 2010;1:216–233.
- Flynn JT, Ng DK, Chan GJ, et al., for the Chronic Kidney Disease in Children Study. The effect of abnormal birth history on ambulatory blood pressure and disease progression in children with chronic kidney disease. *J Pediatr.* 2014;165:154–162.e1.
- Rodriguez MM, Gomez AH, Abitbol CL, et al. Histomorphometric analysis of postnatal glomerulogenesis in extremely preterm infants. *Pediatr Dev Pathol.* 2004;7:17–25.
- Abitbol CL, Bauer CR, Montane B, et al. Long-term follow-up of extremely low birth weight infants with neonatal renal failure. *Pediatr Nephrol.* 2003;18:887–893.
- Hodgin JB, Rasoulpour M, Markowitz GS, D'Agati VD. Very low birth weight is a risk factor for secondary focal segmental glomerulosclerosis. *Clin J Am Soc Nephrol.* 2009;4:71–76.
- Verhave JC, Wetzels JF, van de Kar NC. Novel aspects of atypical haemolytic uraemic syndrome and the role of eculizumab. *Nephrol Dial Transplant.* 2014;29(suppl 4):iv131–iv141.
- Torres VE. Vasopressin receptor antagonists, heart failure, and polycystic kidney disease. *Annu Rev Med.* 2015;66:195–210.
- Jarzembowski T, John E, Panaro F, et al. Impact of non-compliance on outcome after pediatric kidney transplantation: an analysis in racial subgroups. *Pediatr Transplant.* 2004;8:367–371.
- Watson AR. Non-compliance and transfer from paediatric to adult transplant unit. *Pediatr Nephrol.* 2000;14:469–472.
- Aujoulat I, Deccache A, Charles AS, et al. Non-adherence in adolescent transplant recipients: the role of uncertainty in health care providers. *Pediatr Transplant.* 2011;15:148–156.
- Watson AR, Harden PN, Ferris ME, et al. Transition from pediatric to adult renal services: a consensus statement by the International Society of Nephrology (ISN) and the International Pediatric Nephrology Association (IPNA). *Kidney Int.* 2011;80:704–707.
- Watson AR, Harden P, Ferris M, et al. Transition from pediatric to adult renal services: a consensus statement by the International Society of Nephrology (ISN) and the International Pediatric Nephrology Association (IPNA). *Pediatr Nephrol.* 2011;26:1753–1757.
- Gallieni M, Aiello A, Tucci B, et al. The burden of hypertension and kidney disease in Northeast India: the Institute for Indian Mother and Child noncommunicable diseases project. *Scientific World Journal.* 2014;2014:320869.
- White A, Wong W, Sureshkumar P, Singh G. The burden of kidney disease in indigenous children of Australia and New Zealand, epidemiology, antecedent

- factors and progression to chronic kidney disease. *J Paediatr Child Health*. 2010;46:504–509.
33. Zarocostas J. Need to increase focus on non-communicable diseases in global health, says WHO. *BMJ*. 2010;341:c7065.
  34. Gulland A. WHO agrees to set up body to act on non-communicable diseases. *BMJ*. 2013;346:f3483.
  35. Feehally J. Chronic kidney disease: health burden of kidney disease recognized by UN. *Nat Rev Nephrol*. 2011;8:12–13.
  36. Couser WG, Remuzzi G, Mendis S, Tonelli M. The contribution of chronic kidney disease to the global burden of major noncommunicable diseases. *Kidney Int*. 2011;80:1258–1270.

## APPENDIX

Members of the World Kidney Day Steering Committee are Philip Kam Tao Li, Guillermo Garcia-Garcia, William G. Couser, Timur Erk, Julie R Ingelfinger, Kamyar Kalantar-Zadeh, Charles Kernahan, Charlotte Osafo, Miguel C. Riella, Luca Segantini, and Elena Zakharaova. The 4 partners are (in alphabetical order) the IPNA (International Pediatric Nephrology Association), ISN (International Society of Nephrology), ISPD (International Society for Peritoneal Dialysis), and SKCF (Sustainable Kidney Care Foundation).